arXiv:math/0409527v2 [math.OA] 5 Jan 2006

A 1-COHOMOLOGY CHARACTERIZATION OF PROPERTY (T) IN VON NEUMANN ALGEBRAS

JESSE PETERSON

University of California, Los Angeles

Abstract. We obtain a characterization of property (T) for von Neumann algebras in terms of 1-cohomology similar to the Delorme-Guichardet Theorem for groups.

0. Introduction.

The analogue of group representations in von Neumann algebras is the notion of *correspondences* which is due to Connes ([C2], [C3], [P1]), and has been a very useful in defining notions such as property (T) and amenability for von Neumann algebras. It is often useful to view group representations as positive definite functions which we obtain through a GNS construction. Correspondences of a von Neumann algebra N can also be viewed in two separate ways, as *Hilbert* N*-*N *bimodules* $\mathcal{H}$, or as *completely positive maps* $\phi : N \to N$, and the equivalence of these two descriptions is also realized via a GNS construction. This allows one to characterize property (T) for von Neumann algebras in terms of completely positive maps.

For a countable group G there is also a notion of conditionally negative definite functions $\psi : G \to \mathbb{C}$ which satisfy $\psi(g^{-1}) = \overline{\psi(g)}$ and the condition: $\forall n \in \mathbb{N}$, $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{C}$, $g_1, g_2, \dots, g_n \in G$, if $\Sigma_{i=1}^n \alpha_i = 0$ then $\Sigma_{i,j=1}^n \overline{\alpha_j}\alpha_i \psi(g_j^{-1} g_i) \leq 0$. Real valued conditionally negative definite functions can be viewed as cocycles $b \in B^1(G, \pi)$ where $\pi : G \to \mathcal{O}(\mathcal{H})$ is an orthogonal representation of G (see [BdHV]). Real valued conditionally negative definite functions can also be viewed as generators of semigroups of positive definite functions by Schoenberg's Theorem. These equivalences then make it possible for certain connections between 1-cohomology, conditionally negative definite functions, and positive definite deformations, for example the Delorme-Guichardet Theorem [De] [G] which states that a group has property (T) of

---

2000 *Mathematics Subject Classification.* 46L10, 46L57, 22D25.

Key words and phrases. property (T), 1-cohomology, von Neumann algebras.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$-TEX

Kazhdan [Ka] if and only if the first cohomology vanishes for any unitary representation.

It was Evans who introduced the notion of bounded conditionally completely positive/negative maps [E] related to the study the infinitesimal generators of norm continuous semigroups of completely positive maps. He noted that this definition gives an analogue to conditionally positive/negative definite functions on groups. We will extend the notion of conditionally completely negative maps to unbounded maps and use a GNS type construction to alternately view them as closable derivations into a Hilbert N-N bimodule. This is done in the same spirit as ([S1],[S2]) where Sauvageot makes a connection between quantum Dirichlet forms, and differential calculus. Indeed, it is shown in 1.7.1 that conditionally completely negative maps are in fact extensions of generators associated to completely Dirichlet forms, however we are coming from a different perspective here and so we will present the correspondence between conditionally completely negative maps and closable derivations in a way more closely related to group theory.

In studying various properties of groups such as property (T) or the Haagerup property one can give a characterization of these properties in terms of boundedness conditions on conditionally negative definite functions (e.g. [AW]), hence one would hope that this is possible for von Neumann algebras as well.

We will show that one can indeed obtain a characterization of property (T) in this way. The main result is that a separable finite factor has property (T) if and only if the 1-cohomology spaces of closable derivations vanish whenever the domain contains a non-Γ set (see sec. 3 for the definition of a non-Γ set).

0.1. Theorem. *Suppose that N is a separable finite factor, then the following conditions are equivalent:*

(1) N has property (T).

(2) N does not have property Γ and given any weakly dense $*$-subalgebra $N_0 \subset N$, $1 \in N_0$ such that N_0 contains a non-Γ set, we have that every densely defined closable derivation on N_0 into a Hilbert N-N bimodule is inner.

(3) There exists a weakly dense $*$-subalgebra $N_0 \subset N$ such that N_0 is countably generated as a vector space and every closable derivation into a Hilbert N-N bimodule whose domain contains N_0 is inner.

This is the analogue to the Delorme-Guichardet Theorem for groups. As a corollary we obtain that if $X_1, \dots, X_n$ generate a finite factor with property (T), and if at least one of the X_j's has diffuse spectrum then the derivations ∂_{X_i} from [V1] cannot all be closable and hence the conjugate variables cannot all exist in $L^2(N, \tau)$.

0.2. Corollary. *Suppose that N is a finite factor with property (T), let $X_1, \dots, X_n$ generate N as a von Neumann algebra such that $\mathbb{C}[X_1, \dots, X_n]$ contains a non-Γ set. If one of the X_j's has diffuse spectrum then $\Phi^*(X_1, \dots, X_n) = \infty$.*

We also give an application showing that many amalgamated free products of finite von Neumann algebras do not have property (T).

0.3. Theorem. *Let* N_1 *and* N_2 *be finite von Neumann algebras with with normal faithful tracial states* τ_1 *and* τ_2 *respectively, suppose that* B *is a common von Neumann subalgebra such that* $\tau_1|_B = \tau_2|_B$. *If there are unitaries* $u_i \in \mathcal{U}(N_i)$ *such that* $E_B(u_i) = 0$, $i = 1, 2$. *Then* $M = N_1 *_B N_2$ *does not have property (T).*

Other than the introduction there are 4 sections. Section 1 establishes the definitions and notations as well as gives the connection between closable derivations, conditionally completely negative maps, and semigroups of completely positive maps. In section 2 we characterize when a closable derivation is inner in terms of the conditionally completely negative map and the semigroup. In Section 3 we state and prove the main theorem (3.2), and in section 4 we give the application with amalgamated free products (4.2).

1. A GNS-TYPE CONSTRUCTION.

1.1. Conditionally completely negative maps. Let N be a finite von Neumann algebra with normal faithful trace τ.

Definition. *Suppose* $\Psi : N \to L^1(N,\tau)$ *is a* $*$*-preserving linear map whose domain is a weakly dense* $*$*-subalgebra* D_Ψ *of* N *such that* $1 \in D_\Psi$, *then* Ψ *is a conditionally completely negative (c.c.n.) map on* N *if the following condition is satisfied:*

(1.1.1). $\forall n \in \mathbb{N}$, $x_j, y_j \in D_\Psi$, $j \leq n$, *if* $\Sigma_{j=1}^n x_j y_j = 0$ *then* $\Sigma_{i,j=1}^n y_j^* \Psi(x_j^* x_i) y_i \leq 0$.

It is not hard to see that condition (2.1.1) can be replaced with the condition:

(1.1.1)$'$. $\forall n \in \mathbb{N}$, $x_j, y_j \in D_\Psi$, $j \leq n$, if $\Sigma_{j=1}^n x_j y_j = 0$ then $\Sigma_{i,j=1}^n \tau(\Psi(x_j^* x_i) y_i y_j^*) \leq 0$.

If $\phi : N \to N$ is a completely positive map and $k \in N$ then $\Psi(x) = k^* x + xk - \phi(x)$ gives a map which is c.c.n. and bounded. If $\delta : N \to L^2(N,\tau)$ is a derivation then δ is c.c.n. Also if Ψ is a c.c.n. map and $\alpha : N \to N$ is a τ-preserving automorphism then $\Psi' = \alpha \circ \Psi \circ \alpha^{-1}$ is another c.c.n. map.

One can check that if Ψ_1 and Ψ_2 are c.c.n. such that $D_{\Psi_1} \cap D_{\Psi_2}$ is weakly dense in N, and if $s, t \geq 0$, then $\Psi = s\Psi_1 + t\Psi_2$ is c.c.n. Also if $\{\Psi_t\}_t$ is a family of c.c.n. maps on the same domain and Ψ is the pointwise $\| \cdot \|_1$-limit of $\{\Psi_t\}_t$ then Ψ is c.c.n.

We say that Ψ is symmetric if $\tau(\Psi(x)y) = \tau(x\Psi(y)), \forall x, y \in D_\Psi$. We say that Ψ is conservative if $\tau \circ \Psi = 0$. We also say that Ψ is closable if the quadratic form q on $L^2(N,\tau)$ given by $D(q) = D_\Psi$, $q(x) = \tau(\Psi(x)x^*)$ is closable. Note that we will see in (1.3) that if $\Psi : D_\Psi \to L^2(N,\tau) \subset L^1(N,\tau)$ is a conservative symmetric c.c.n. map then Ψ is automatically closable.

Note that if Ψ is a conservative symmetric c.c.n. map then $\tau(\Psi(1)x) = \tau(\Psi(x)) = 0, \forall x \in D_\Psi$, hence $\Psi(1) = 0$. Also note that if Ψ is symmetric and $\Psi(1) \geq 0$ then given any $x \in D_\Psi$, if we let $x_1 = x, x_2 = 1, y_1 = -1, y_2 = x$, then the above condition implies that $\tau(\Psi(x)x^*) \geq 0$, so that we actually have positivity instead of just the symmetry condition.

1.2. Closable derivations. Let $\mathcal{H}$ be a Hilbert N-N bimodule, a derivation of N is a (possibly unbounded) map $\delta : N \to \mathcal{H}$ which is defined on a weakly dense $*$-subalgebra D_δ of N such that $1 \in D_\delta$, and such that $\forall x, y \in D_\delta$, $\delta(xy) = x\delta(y) + \delta(x)y$. δ is closable if it is closable as an operator from $L^2(N, \tau)$ to $\mathcal{H}$.

δ is inner if $\delta(x) = x\xi - \xi x$ for some $\xi \in \mathcal{H}$. δ is spanning if $\overline{sp} D_\delta \delta(D_\delta) = \mathcal{H}$. δ is real if $\langle x\delta(y), \delta(z)\rangle_{\mathcal{H}} = \langle \delta(z^*), \delta(y^*)x^*\rangle_{\mathcal{H}}$, $\forall x, y, z \in D_\delta$.

If $\delta' : D_\delta \to \mathcal{H}'$ is another derivation then we say that δ and δ' are equivalent if there exists a unitary map $U : \mathcal{H} \to \mathcal{H}'$ such that $U(x\delta(y)z) = xU(\delta(y))z = x\delta'(y)z$ for all $x, y, z \in D_\delta$.

Recall that if $\mathcal{H}$ is a Hilbert N-N bimodule then we can define the adjoint bimodule $\mathcal{H}^o$ where $\mathcal{H}^o$ is the conjugate Hilbert space of $\mathcal{H}$ and the bimodule structure is given by $x\xi^o y = (y^*\xi x^*)^o$. If $\delta : D_\delta \to \mathcal{H}$ is a closable derivation then we may define the adjoint derivation $\delta^o : D_\delta \to \mathcal{H}^o$ by setting $\delta^o(x) = \delta(x^*)^o$, then δ^o is a closable derivation and furthermore the derivations $\frac{1}{2}(\delta + \delta^o)$, and $\frac{1}{2}(\delta - \delta^o)$ are real derivations from D_δ to $\mathcal{H} \oplus \mathcal{H}^o$.

1.3. From conditionally completely negative maps to closable derivations. Let Ψ be a conservative symmetric c.c.n. map on N with domain D_Ψ. We associate to Ψ a derivation in the following way (compare with [S1]):

let $\mathcal{H}_0 = \{\Sigma_{i=1}^n x_i \otimes y_i \in D_\Psi \otimes D_\Psi | \Sigma_{i=1}^n x_i y_i = 0\}$. Define a sesquilinear form on $\mathcal{H}_0$ by $\langle \Sigma_{i=1}^n x_i' \otimes y_i', \Sigma_{j=1}^m x_j \otimes y_j\rangle_\Psi = -\frac{1}{2}\Sigma_{i=1}^n \Sigma_{j=1}^m \tau(\Psi(x_j^* x_i') y_i' y_j^*)$. The positivity of $\langle \cdot, \cdot \rangle_\Psi$ is equivalent to the c.c.n. condition on Ψ. Let $\mathcal{H}$ be the closure of $\mathcal{H}_0$ after we mod out by the kernel of $\langle \cdot, \cdot \rangle_\Psi$. If $p = \Sigma_{k=1}^n x_k \otimes y_k$ such that $\Sigma_{k=1}^n x_k y_k = 0$ then $x \mapsto -\frac{1}{2}\Sigma_{i,j=1}^n \tau(x_j^* x x_i \Psi(y_i y_j^*))$ is a positive normal functional on N with norm $\langle p, p\rangle_\Psi$. Similarly $y \mapsto -\frac{1}{2}\Sigma_{i,j=1}^n \tau(\Psi(x_j^* x_i) y_i y y_j^*)$ is a positive normal functional on N with norm $\langle p, p\rangle_\Psi$. We also have left and right commuting actions of D_Ψ on $\mathcal{H}_0$ given by $xpy = x(\Sigma_{k=1}^n x_k \otimes y_k)y = \Sigma_{k=1}^n (xx_k) \otimes (y_k y)$, and by the preceeding remarks we have $\langle xp, xp\rangle_\Psi = \langle x^*xp, p\rangle_\Psi \leq \|x^*x\| \langle p, p\rangle_\Psi = \|x\|^2 \langle p, p\rangle_\Psi$ and $\langle py, py\rangle_\Psi \leq \|y\|^2 \langle p, p\rangle_\Psi$ for all $x, y \in D_\Psi$. Hence the above actions of D_Ψ pass to commuting left and right actions on $\mathcal{H}$, and they extend to left and right actions of N on $\mathcal{H}$ given by the formulas:

$$\langle x[\Sigma_{i=1}^n x_i' \otimes y_i'], [\Sigma_{j=1}^m x_j \otimes y_j]\rangle_{\mathcal{H}} = \Sigma_{i=1}^n \Sigma_{j=1}^m \tau(x_j^* x x_i' \Psi(y_i' y_j^*)),$$

$$\langle [\Sigma_{i=1}^n x_i' \otimes y_i'] y, [\Sigma_{j=1}^m x_j \otimes y_j]\rangle_{\mathcal{H}} = \Sigma_{i=1}^n \Sigma_{j=1}^m \tau(\Psi(x_j^* x_i') y_i' y y_j^*).$$

Since the above forms are normal the left and right actions commute and are normal thus making $\mathcal{H}$ into a Hilbert N-N bimodule.

Define $\delta_\Psi : D_\Psi \to \mathcal{H}$ to be given by $\delta_\Psi(x) = [x \otimes 1 - 1 \otimes x]$. Then δ_Ψ is a derivation such that

$$\langle \delta_\Psi(x), \delta_\Psi(y)\rangle_{\mathcal{H}} = \langle x \otimes 1 - 1 \otimes x, y \otimes 1 - 1 \otimes y\rangle_\Psi$$

$$= -\frac{1}{2}\tau(\Psi(y^*x)) + \frac{1}{2}\tau(\Psi(x)y^*) + \frac{1}{2}\tau(\Psi(y^*)x) - \frac{1}{2}\tau(\Psi(1)xy^*)$$
$$= \tau(\Psi(x)y^*),$$

for all $x, y \in D_\Psi$. Also δ_Ψ is real since

$$\langle x\delta_\Psi(y), \delta_\Psi(z)\rangle_{\mathcal{H}} = \langle xy \otimes 1 - x \otimes y, z \otimes 1 - 1 \otimes z\rangle_\Psi$$

$$= -\frac{1}{2}\tau(\Psi(z^*xy)) + \frac{1}{2}\tau(\Psi(xy)z^*) + \frac{1}{2}\tau(\Psi(z^*x)y) - \frac{1}{2}\tau(\Psi(x)yz^*)$$
$$= -\frac{1}{2}\tau(\Psi(1)z^*xy) + \frac{1}{2}\tau(\Psi(z^*)xy) + \frac{1}{2}\tau(\Psi(y)z^*x) - \frac{1}{2}\tau(\Psi(yz^*)x)$$
$$= \langle 1 \otimes z^* - z^* \otimes 1, 1 \otimes y^*x^* - y^* \otimes x^*\rangle_\Psi = \langle \delta_\Psi(z^*), \delta_\Psi(y^*)x^*\rangle_{\mathcal{H}},$$

for all $x, y, z \in D_\Psi$. If Ψ is closable then it follows that δ_Ψ is closable. Also note that if $\Psi : D_\Psi \to L^2(N, \tau) \subset L^1(N, \tau)$ then we would have that $D_\Psi = D(\delta_\Psi^* \delta_\Psi)$ which would show that δ (and hence also Ψ) is closable.

Note that we will assume in addition that δ_Ψ is spanning by restricting ourselves to $\overline{sp} D_\Psi \delta(D_\Psi) \subset \mathcal{H}$.

Also note that the requirement that $\Psi(1) = 0$ is not really much of a restriction since if Ψ is any symmetric c.c.n. map with $\Psi(1) \in L^2(N, \tau)$ then $\Psi'(x) = \Psi(x) - \frac{1}{2}\Psi(1)x - \frac{1}{2}x\Psi(1)$ defines a symmetric c.c.n. map with $\Psi'(1) = 0$.

1.4. From closable derivations to conditionally completely negative maps. Let $\mathcal{H}$ be a Hilbert N-N bimodule and suppose that $\delta : N \to \mathcal{H}$ is a closable real derivation defined on a weakly dense $*$-subalgebra D_δ of N with $1 \in D_\delta$.

Let $D_\Psi = \{x \in D(\overline{\delta}) \cap N | y \mapsto \langle \overline{\delta}(x), \delta(y^*)\rangle$ gives a normal linear functional on $N\}$ then by [S2] and [DL] $D(\overline{\delta}) \cap N$ is a $*$-subalgebra and hence one can show that D_Ψ is a $*$-subalgebra of N. We define the map $\Psi_\delta : D_\Psi \to L^1(N, \tau)$ by letting $\Psi_\delta(x)$ be the Radon-Nikodym derivative of the normal linear functional $y \mapsto \langle \overline{\delta}(x), \delta(y^*)\rangle$. Since δ is closable Ψ_δ is also closable.

As δ is real Ψ_δ is a symmetric $*$-preserving map such that $\tau \circ \Psi = 0$ and if $n \in \mathbb{N}$, $x_1, x_2, \ldots, x_n, y_1, y_x, \ldots, y_n \in D_\Psi$, such that $\Sigma_{i=1}^n x_i y_i = 0$ then:

$$\Sigma_{i,j=1}^n \tau(\Psi(x_j^* x_i) y_i y_j^*) = \Sigma_{i,j=1}^n \langle \delta(x_j^* x_i), \delta(y_j y_i^*)\rangle_{\mathcal{H}}$$

$$= \Sigma_{i,j=1}^n \langle x_j^* \delta(x_i), y_j \delta(y_i^*) + \delta(y_j) y_i^*\rangle_{\mathcal{H}} + \langle \delta(x_j^*) x_i, y_j \delta(y_i^*) + \delta(y_j) y_i^*\rangle_{\mathcal{H}}$$
$$= \Sigma_{i,j=1}^n \langle \delta(x_i) y_i, x_j \delta(y_j)\rangle_{\mathcal{H}} + \langle x_i \delta(y_i), \delta(x_j) y_j\rangle_{\mathcal{H}}$$
$$= -2\|\Sigma_{i=1}^n \delta(x_i) y_i\|_{\mathcal{H}}^2 \leq 0.$$

Hence Ψ_δ is a conservative symmetric c.c.n. map on D_Ψ.

Note that if we restrict ourselves to closable derivations which are spanning then an easy calculation shows that the constructions above are inverses of each other in the sense that $\Psi_{\delta_\Psi}|_{D_\Psi} = \Psi$ and $\overline{\delta_{\Psi_\delta}} \cong \overline{\delta}$.

1.5. Closable derivations and c.c.n. maps from groups. Let Γ be a discrete group, (C, τ_0) a finite von Neumann algebra with a normal faithful trace, and σ a cocycle action of Γ on (C, τ_0) by τ_0-preserving automorphisms. Denote by $N = C \rtimes_\sigma \Gamma$ the corresponding cross-product algebra with trace τ given by $\tau(\Sigma c_g u_g) = \tau_0(c_e)$, where $c_g \in C$ and $\{u_g\}_g \subset N$ denote the canonical unitaries implementing the action σ on C.

Let $(\pi_0, \mathcal{H}_0)$ be a unitary or orthogonal representation of Γ, and let $b : \Gamma \to \mathcal{H}_0$ be an (additive) cocycle of Γ, i.e. $b(gh) = \pi_0(g)b(h) + b(g)$, $\forall g, h \in \Gamma$. Set $\mathcal{H}_{\pi_0}$ to be the Hilbert space $\mathcal{H}_0 \overline{\otimes}_{\mathbb{R}} L^2(N, \tau)$ if π_0 is an orthogonal representation and $\mathcal{H}_0 \overline{\otimes}_{\mathbb{C}} L^2(N, \tau)$ if π_0 is a unitary representation. We let N act on the right of $\mathcal{H}_{\pi_0}$ by $(\xi \otimes \hat{x})y = \xi \otimes (\widehat{xy})$, $x, y \in N, \xi \in \mathcal{H}_0$ and on the left by $c(\xi \otimes \hat{x}) = \xi \otimes (\widehat{cx}), u_g(\xi \otimes \hat{x}) = (\pi_0(g)\xi) \otimes (\widehat{u_g x})$, $c \in C, x \in N, g \in \Gamma, \xi \in \mathcal{H}_0$. Let D_Γ be the $*$-subalgebra generated by C and $\{u_g\}_g$, we define δ_b by $\delta_b(c_g u_g) = c_g \delta_b(u_g) = b(g) \otimes \widehat{c_g u_g}$, $c_g \in C, g \in \Gamma$, then we can extend δ_b linearly so that δ_b is a derivation on D_Γ. If $(\pi_0, \mathcal{H}_0)$ is an orthogonal representation and 1_g denotes the Dirac delta function at g then:

$$\langle c u_g \delta_b(u_h), \delta_b(u_k)\rangle = \langle \pi_0(g)b(h), b(k)\rangle \langle \widehat{c u_g u_h}, \hat{u}_k\rangle$$

$$= \langle -\pi_0(g)\pi_0(h)b(h^{-1}), -\pi_0(k)b(k^{-1})\rangle \langle \widehat{c u_g u_h}, \hat{u}_k\rangle 1_k(gh)$$

$$= \langle b(k^{-1}), b(h^{-1})\rangle \langle \hat{u}_k^*, \widehat{u_h^* u_g^* c^*}\rangle$$

$$= \langle \delta_b(u_k^*), \delta_b(u_h^*)u_g^* c^*\rangle,$$

for all $g, h, k \in \Gamma$, $c \in C$, thus showing that δ_b is real.

Also we have:

$$|\langle \delta_b(c_g u_g), \delta_b(\Sigma_{h \in \Gamma} d_h u_h)\rangle| = |\Sigma_{h \in \Gamma} \langle b(g), b(h)\rangle \langle \widehat{c_g u_g}, \widehat{d_h u_h}\rangle|$$

$$= \|b(g)\|^2 |\langle \widehat{c_g u_g}, \Sigma_{h \in \Gamma} \widehat{d_h u_h}\rangle| \leq \|b(g)\|^2 \|c_g\| \|\Sigma_{h \in \Gamma} d_h u_h\|_1,$$

for all $g \in \Gamma$, $c_g \in C$, $\Sigma_{h \in \Gamma} d_h u_h \in D_\Gamma$. Hence if $x = \Sigma_{g \in \Gamma} c_g u_g \in D_\Gamma$, and $y \in D_\Gamma$ then $|\langle \delta_b(x), \delta_b(y)\rangle| \leq (\Sigma_{g \in \Gamma} \|b(g)\|^2 \|c_g\|)\|y\|_1$. In particular this shows that δ_b is closable.

Now suppose that $\psi : \Gamma \to \mathbb{C}$ is a real valued conditionally negative definite function on Γ such that $\psi(e) = 0$ and let (π_ψ, b_π) be the representation and cocycle which correspond to ψ through the GNS construction [BdHV]. Let $(\mathcal{H}, \delta)$ denote the Hilbert N-N bimodule and closable derivation constructed out of (π_ψ, b_π) as above and let Ψ be the symmetric c.c.n. map associated to $(\mathcal{H}, \delta)$ as in 1.4. Then a calculation shows that $\Psi(\Sigma_g c_g u_g) = \Sigma_g \psi(g) c_g u_g$, and in fact it is an easy exercise to show that even if ψ is not real valued $\Psi(\Sigma_g c_g u_g) = \Sigma_g \psi(g) c_g u_g$ still describes a c.c.n. map.

Conversely, if $(\mathcal{H}, \delta)$ is a Hilbert N-N bimodule and a closable derivation such that δ is defined on the $*$-subalgebra generated by C and $\{u_g\}_g$ then we can associate to it a representation π_0 on $\mathcal{H}_0 = \overline{sp}\{\delta(u_g)u_g^* | g \in \Gamma\}$ by $\pi_0(g)\xi' = u_g \xi' u_g^*$, $\xi' \in \mathcal{H}_0$. Also we may associate to δ a group cocycle b on Γ by $b(g) = \delta(u_g)u_g^*$, $g \in \Gamma$. If Ψ is a c.c.n. map which is also defined on the $*$-subalgebra generated by C and $\{u_g\}_g$ then we can associate to it a conditionally negative definite function ψ by $\psi(g) = \tau(\Psi(u_g)u_g^*)$. Furthermore if δ is real then by taking only the real span above we have that $\mathcal{H}_0$ is a real Hilbert space and π_0 is an orthogonal representation, also ψ is real valued if and only if Ψ is symmetric, and if $(\mathcal{H}, \delta)$ and Ψ correspond to each other as in 1.3 and 1.4 then (π_0, b) and ψ correspond to each other via the GNS construction.

1.6. Examples from free probability. From above we have two main examples of closable derivations, those which are inner, and those which come from cocycles on groups. In [V1] and [V2] Voiculescu uses certain derivations in a key role for his non-microstates approach to free entropy and mutual free information. We will recall these derivations which will give us more examples of closable derivations under certain circumstances.

1.6.1 The derivation ∂_X from [V1]. Suppose $B \subset N$ is a $*$-subalgebra with $1 \in B$ and $X = X^* \in N$. If we denote by $B[X]$ the subalgebra generated by B and X, and if X and B are algebraically free (i.e. they do not satisfy any nontrivial algebraic relations) then there is a well-defined unique derivation

$$\partial_X : B[X] \to B[X] \otimes B[X] \subset L^2(N, \tau) \otimes L^2(N, \tau)$$

such that $\partial_X(X) = 1 \otimes 1$ and $\partial_X(b) = 0 \ \forall b \in B$.

We note that if ∂_X is inner then by identifying $L^2(N, \tau) \otimes L^2(N, \tau)$ with the Hilbert-Schmidt operators we would have that there exists a Hilbert-Schmidt operator which commutes with B. Therefore if B contains a diffuse element (i.e. an element which generates a von Neumann algebra without minimal projections) then we must have that ∂_X is not inner.

Recall from [V1] that the conjugate variable $J(X : B)$ of X w.r.t. B is an element in $L^1(W^*(B[X]), \tau)$ such that $\tau(J(X : B)m) = \tau \otimes \tau(\partial_X(m)) \ \forall m \in B[X]$, i.e. $J(X : B) = \partial_X^*(1 \otimes 1)$.

If $J(X : B)$ exists and is in $L^2(N, \tau)$ (as in the case when we perturb a set of generators by free semicircular elements) then by Corollary 4.2 in [V1] we have that ∂_X is a closable derivation.

1.6.2 The derivation $\delta_{A:B}$ from [V2]. Suppose $A, B \subset N$ are two $*$-subalgebras with $1 \in A, B$. If we denote by $A \vee B$ the subalgebra generated by A and B, and if A and B are algebraically free then we may define a unique derivation

$$\delta_{A:B} : A \vee B \to (A \vee B) \otimes (A \vee B) \subset L^2(N, \tau) \otimes L^2(N, \tau)$$

by $\delta_{A:B}(a) = a \otimes 1 - 1 \otimes a \ \forall a \in A$, and $\delta_{A:B}(b) = 0 \ \forall b \in B$.

We note that for the same reason as above if B contains a diffuse element and $A \neq \mathbb{C}$ then we must have that the derivation is not inner.

Recall from [V2] that the liberation gradient $j(A : B)$ of (A, B) is an element in $L^1(W^*(A \cup B), \tau)$ such that $\tau(j(A : B)m) = \tau \otimes \tau(\delta_{A:B}(m)) \ \forall m \in A \vee B$, i.e. $j(A : B) = \delta_{A:B}^*(1 \otimes 1)$.

If $j(A : B)$ exists and is in $L^2(N, \tau)$ then by Corollary 6.3 in [V2] $\delta_{A:B}$ is a closable derivation.

1.7. Generators of completely positive semigroups. Let N be a finite von Neumann algebra with normal faithful trace τ. A weak*-continuous semigroup $\{\phi_t\}_{t\geq 0}$ on N is said to be symmetric if $\tau(x\phi_t(y)) = \tau(\phi_t(x)y)$, $\forall x, y \in N$, and completely Markovian if each ϕ_t is a unital c.p. map on N. We denote by Δ the generator of a symmetric completely Markovian semigroup $\{\phi_t\}_{t\geq 0}$ on N, i.e. Δ is the densely defined operator on N described by $D(\Delta) = \{x \in N : \frac{x-\phi_t(x)}{t} \text{ has a weak limit as } t \to 0\}$, and $\Delta(x) = \lim_{t\to 0} \frac{x-\phi_t(x)}{t}$, we also denote by Δ the generator of the corresponding semigroup on $L^2(N, \tau)$. Then Δ describes a completely Dirichlet form [DL] on $L^2(N, \tau)$ by $D(\mathcal{E}) = D(\Delta^{1/2})$, $\mathcal{E}(x) = \|\Delta^{1/2}(x)\|_2^2$.

From [DL] we have that $D(\mathcal{E}) \cap N$ is a weakly dense $*$-subalgebra and hence it follows from [S1] that there exists a Hilbert N-N bimodule $\mathcal{H}$ and a closeable derivation $\delta : D(\mathcal{E}) \cap N \to \mathcal{H}$ such that $\mathcal{E}(x) = \|\delta(x)\|^2$, $\forall x \in D(\mathcal{E}) \cap N$. Conversely it follows from [S2] that if $D(\delta)$ is a weakly dense $*$-subalgebra with $1 \in D(\delta)$ and $\delta : D(\delta) \to \mathcal{H}$ is a closable derivation, then the closure of the quadratic form given by $\|\delta(x)\|^2$ is completely Dirichlet on $L^2(N, \tau)$ and hence generates a symmetric completely Markovian semigroup as above (see also [CiSa]).

From sections 1.3 and 1.4, and from the remarks above we obtain the following.

1.7.1. Theorem. *Let $N_0 \subset N$ be a weakly dense $*$-subalgebra with $1 \in N_0$ and suppose $\Psi : N_0 \to L^1(N, \tau)$ is a closable, conservative, symmetric c.c.n. map, such that $\Psi^{-1}(L^2(N, \tau))$ is weakly dense in N. Then $\Delta = \Psi|_{\Psi^{-1}(L^2(N,\tau))}$ is closable as a densely defined operator on $L^2(N, \tau)$ and $\overline{\Delta}$ is the generator of a symmetric completely Markovian semigroup on N. Conversely if Δ is the generator of a symmetric completely Markovian semigroup on N then Δ extends to a conservative, symmetric c.c.n. map $\Psi : N_0 \to L^1(N, \tau)$ where N_0 is the $*$-subalgebra generated by $D(\Delta)$.*

2. A CHARACTERIZATION OF INNER DERIVATIONS.

Let N be a finite von Neumann algebra with normal faithful trace τ. Given a symmetric c.c.n. map Ψ on N we will now give a characterization of when Ψ is norm bounded.

2.1. Theorem. *Let $\Psi : D_\Psi \to L^1(N, \tau)$ be a closable, conservative, symmetric c.c.n. map with weakly dense domain D_Ψ. Let $\delta : D_\Psi \to \mathcal{H}$ be the closable derivation associated with Ψ. Then the following conditions are equivalent:*

(a) δ extends to an everywhere defined derivation δ' which is inner and such that given any $x \in N$ there exists a constant $C_x > 0$ such that $|\langle \delta'(x), \delta'(y)\rangle| \leq C_x \|y\|_1$, $\forall y \in N$.

(b) There exists a constant $C > 0$ such that $|\langle \delta(x), \delta(y)\rangle| \leq C\|x\|\|y\|_1$, $\forall x, y \in D_\Psi$.

(c) Ψ is norm bounded on $(D_\Psi)_1$.

(d) The image of Ψ is contained in $N \subset L^1(N, \tau)$ and $-\Psi$ extends to a mapping which generates a norm continuous semigroup of normal c.p. maps.

(e) There exists $k \in N$ and a normal c.p. map $\phi : N \to N$ such that $\Psi(x) = k^*x + xk - \phi(x)$, $\forall x \in D_\Psi$.

Proof. (a) ⇒ (c): Let δ' be the everywhere defined extension of δ, and let Ψ' be the c.c.n. map associated with δ'. Since given any $x \in N$ there exists a constant $C_x > 0$ such that $|\langle \delta'(x), \delta'(y)\rangle| \leq C_x\|y\|_1$, $\forall y \in N$ we have that the image of Ψ' is contained in N. Also since $\Psi'(1) = 0$ we have that for all $x \in D_{\Psi'}$, $\Psi'(x^*x) - x^*\Psi'(x) - \Psi'(x^*)x \leq 0$ and so $-\Psi'$ is a dissipation ([L],[Ki]). As $-\Psi'$ is also everywhere defined, it is bounded by Theorem 1 in [Ki].

(b) ⇔ (c): Suppose (b) holds, then for all $x, y \in D_\Psi$,

$$|\tau(\Psi(x)y^*)| = |\langle \delta(x), \delta(y)\rangle| \leq C\|x\|\|y\|_1.$$

So by taking the supremum over all $y \in D_\Psi$ such that $\|y\|_1 \leq 1$ we have that $\|\Psi(x)\| \leq C\|x\|$, $\forall x \in D_\Psi$.

Suppose now that Ψ is bounded by $C > 0$. Then for all $x, y \in D_\Psi$,

$$|\langle \delta(x), \delta(y)\rangle| = |\tau(\Psi(x)y^*)| \leq \|\Psi(x)\|\|y\|_1 \leq C\|x\|\|y\|_1.$$

(c) ⇒ (d): This follows from [E] Proposition 2.10.

(d) ⇒ (e): This is Theorem 3.1 in [ChE].

(e) ⇒ (a): Suppose that for $k \in N$ and ϕ c.p. we have $\Psi(x) = k^*x + xk - \phi(x)$, $\forall x \in N$. Let $\phi' = \tau(\phi(1))^{-1}\phi$ and let $(\mathcal{H}, \xi)$ be the pointed Hilbert N-bimodule associated with ϕ'. Hence if we set $\delta'(x) = (\tau(\phi(1))/2)^{1/2}[x, \xi]$ then we have $\delta' \cong \delta$. By replacing k with $\frac{1}{2}(k + k^*)$ and ϕ with $\frac{1}{2}(\phi + \phi^*)$ we may assume that ϕ is symmetric, it is then easy to verify that there exists a constant $C > 0$ such that for all $x, y \in N$, $|\langle \delta'(x), \delta'(y)\rangle| \leq C\|x\|\|y\|_1$. Hence δ' gives an everywhere defined extension of δ which satisfies the required properties. □

Our next result is in the same spirit as Theorem 2.1 and provides several equivalent conditions for when a closable derivation is inner.

2.2. Theorem. *Let $\Psi : D_\Psi \to L^1(N,\tau)$ be a closable, conservative, symmetric c.c.n. map with weakly dense domain D_Ψ. Let $\delta : D_\Psi \to \mathcal{H}$ be the closable derivation associated with Ψ. Then the following conditions are equivalent:*

(α) δ is inner.

(β) δ is bounded on $(D_\Psi)_1$.

(γ) Ψ is $\| \cdot \|_1$-bounded on $(D_\Psi)_1$.

(δ) Ψ can be approximated uniformly by c.p. maps in the following sense: for all $\varepsilon > 0$, there exists $k \in N$, and ϕ a normal c.p. map such that $\|\Psi(x) - k^*x - xk + \phi(x)\|_1 \leq \varepsilon \|x\|$, $\forall x \in D_\Psi$.

Proof. (α) $\Rightarrow$ (δ): Suppose $\xi \in \mathcal{H}$ such that $\delta(x) = x\xi - \xi x$, $\forall x \in D_\Psi$. Let $\varepsilon > 0$. Since the subspace of "left and right bounded" vectors is dense in $\mathcal{H}$, let $\xi_0 \in \mathcal{H}$ such that there exists a constant $C > 0$ such that $\|x\xi_0\| \leq C\|x\|_2$, $\forall x \in N$, $\|\xi_0\| \leq \|\xi\|$, and also $\|\xi - \xi_0\| < \varepsilon / 8\|\xi\|$. As ξ_0 is "bounded" we may let ϕ_{ξ_0} be the normal c.p. map associated with $\xi_0/\|\xi_0\|$. Let $\phi = 2\|\xi_0\|^2\phi_{\xi_0}$, and let $k = \phi(1)/2$.

Note that since δ is real we have that ξ_0 is also real, i.e. $\langle x\xi_0, \xi_0 y\rangle = \langle y^*\xi_0, \xi_0 x^*\rangle$, $\forall x, y \in N$.

Then if $x, y \in D_\Psi$ we have:

$$\tau((\Psi(x) - k^*x - xk + \phi(x))y^*)$$

$$\begin{aligned} &= \tau(\Psi(x)y^*) - \frac{1}{2}\tau(\phi(1)xy^*) - \frac{1}{2}\tau(x\phi(1)y^*) + \tau(\phi(x)y^*) \\ &= \langle \delta(x), \delta(y)\rangle - \langle xy^*\xi_0, \xi_0\rangle - \langle y^*x\xi_0, \xi_0\rangle + 2\langle x\xi_0 y^*, \xi_0\rangle \\ &= \langle x\xi - \xi x, y\xi - \xi y\rangle - \langle x\xi_0 - \xi_0 x, y\xi_0 - \xi_0 y\rangle. \end{aligned}$$

Hence:

$$|\langle \Psi(x) - k^*x - xk + \phi(x), y\rangle|$$

$$\begin{aligned} &\leq \|x\xi - \xi x\|\|y\xi - \xi y - y\xi_0 + \xi_0 y\| + \|y\xi_0 - \xi_0 y\|\|x\xi - \xi x - x\xi_0 + \xi_0 x\| \\ &\leq 4\|x\|\|\xi\|\|y\|\|\xi - \xi_0\| + 4\|y\|\|\xi_0\|\|x\|\|\xi - \xi_0\| \\ &\leq \varepsilon\|x\|\|y\|. \end{aligned}$$

Thus by taking the supremum over all $y \in (D_\Psi)_1$ we have the desired result.

(δ) $\Rightarrow$ (γ): Let $k \in N$ and ϕ c.p. such that $\|\Psi(x) - k^*x - xk + \phi(x)\|_1 \leq \|x\|$. By ([P2] 1.1.2) $\|\phi(x)\|_2 \leq \|\phi(1)\|_2\|x\|$, $\forall x \in N$. Hence for all $x \in D_\Psi$:

$$\|\Psi(x)\|_1 \leq \|\Psi(x) - k^*x - xk + \phi(x)\|_1 + \|k^*x - xk + \phi(x)\|_2$$

$$\leq (1 + 2\|k\|_2 + \|\phi(1)\|_2)\|x\|.$$

Thus Ψ is bounded in $\|\cdot\|_1$ on $(D_\Psi)_1$.

$(\gamma) \Rightarrow (\beta)$: Suppose $\|\Psi(x)\|_1 \leq C\|x\|$, $\forall x \in D_\Psi$, then:

$$\|\delta(x)\|^2 = \tau(\Psi(x)x^*) \leq \|\Psi(x)\|_1\|x\| \leq C\|x\|^2,$$

for all $x \in D_\Psi$.

$(\beta) \Rightarrow (\alpha)$: As δ is bounded on $(D_\Psi)_1$ we may extend δ to a derivation on the C^*-algebra A which is generated by D_Ψ. Let $X = \{\delta(u)u^* | u \in \mathcal{U}(A)\}$, for each $v \in \mathcal{U}(A)$ we let v act on $\mathcal{H}$ by $v \cdot \xi = v\xi + \delta(v)$. Let ξ_0 be the center of the set X. Then since $\forall \xi, \eta \in \mathcal{H}$, $\|v \cdot \xi - v \cdot \eta\| = \|\xi - \eta\| = \|\xi v - \eta v\|$, the center of the set $v \cdot X$ is $v \cdot \xi_0$, and the center of the set Xv is $\xi_0 v$. Further we have that $v \cdot X = Xv$ and thus $v \cdot \xi_0 = \xi_0 v$. Since v was arbitrary and every $x \in A$ is a linear combination of unitaries we have that $\delta(x) = \xi_0 x - x\xi_0$, $\forall x \in A$. ☐

In general we may have that Ψ is unbounded in $\|\cdot\|_1$ even if $\|\phi_t(x) - x\|_2$ converges to 0 uniformly on N_1. However we will show in the next section that if N has property (T) and the domain of Ψ contains a "critical set" as in Proposition 1 of [CJ] then this cannot happen.

3. Property (T) in terms of closable derivations.

Given a finite von Neumann algebra M with countable decomposable center. We will say that M has property (T) if the inclusion $(M \subset M)$ is rigid in the sense of [P2], i.e. M has property (T) if and only if there exists a normal faithful tracial state τ' on M such that one of the following equivalent conditions hold:

1. $\forall \varepsilon > 0$, $\exists F' \subset M$ finite and $\delta' > 0$ such that if $\mathcal{H}$ is a Hilbert $M-M$ bimodule with a vector $\xi \in \mathcal{H}$ satisfying the conditions $\|\langle \cdot \xi, \xi\rangle - \tau'\| \leq \delta'$, $\|\langle \xi \cdot, \xi\rangle - \tau'\| \leq \delta'$, and $\|y\xi - \xi y\| \leq \delta'$, $\forall y \in F'$ then $\exists \xi_0 \in \mathcal{H}$ such that $\|\xi_0 - \xi\| \leq \varepsilon$ and $x\xi_0 = \xi_0 x$, $\forall x \in M$.
2. $\forall \varepsilon > 0$, $\exists F \subset M$ finite and $\delta > 0$ such that if $\phi : M \to M$ is a normal, completely positive map with $\tau' \circ \phi \leq \tau'$, $\phi(1) \leq 1$ and $\|\phi(y) - y\|_2 \leq \delta$, $\forall y \in F$, then $\|\phi(x) - x\|_2 \leq \varepsilon$, $\forall x \in M$, $\|x\| \leq 1$.

Furthermore it was shown in [P2] that the above definition is independent of the trace τ', and in the case when N is a factor this agrees with the original definition in [CJ].

In this section we will obtain a characterization of property (T) in terms of certain boundedness conditions on c.c.n. maps. As we are dealing with unbounded maps the domain of a map will be of crucial importance. We will thus want to consider c.c.n. maps whose domain contains a "critical set", which by Remark 4.1.6 of [P1] motivates the following.

3.1. Definition. *Suppose that N is a II_1 factor and let $N_0 \subset N$ be a weakly dense $*$-subalgebra of N, $1 \in N_0$, then N_0 contains a non-Γ set if $\exists F \subset N_0$ finite, $K > 0$ such that $\forall \xi \in L^2(N, \tau)$, if $\langle \xi, 1 \rangle = 0$ then we have $\|\xi\|_2^2 \leq K\Sigma_{x\in F}\|x\xi - \xi x\|_2^2$.*

Note that by Lemma 2.4 of [C1] one can check that N_0 has a non-Γ set if and only if there exists a finitely generated subgroup $\mathcal{G} \subset \mathrm{Int}\, C^*(N_0)$ such that there is no non-normal $\mathcal{G}$-invariant state on N. Also it follows from the definition that $N \subset N$ contains a non-Γ set if and only if N does not have property Γ of Murray and von Neumann [MvN]. Also note that if Λ is a countable ICC group then by [Ef] Λ is not inner amenable if and only if $\mathbb{C}\Lambda$ contains a non-Γ set.

We now come to the main result which is to give several equivalent characterizations of property (T), in particular we obtain a 1-cohomology characterization of property (T) which is the analogue of the Delorme-Guichardet Theorem from group theory.

3.2. Theorem. *Suppose that N is a separable finite factor with normal faithful trace τ. Let $N_0 \subset N$ be a weakly dense $*$-subalgebra such that $1 \in N_0$ and N_0 is countably generated as a vector space. Consider the following conditions:*

(a) N has property (T).

(b) $\exists F \subset N_0$ finite and $K > 0$ such that if $\mathcal{H}$ is a Hilbert N-N bimodule, $\xi \in \mathcal{H}$, and if $\delta_\xi = \max_{x\in F}\{\|x\xi - \xi x\|\}$ then $\exists \xi_0 \in \mathcal{H}$ such that $x\xi_0 = \xi_0 x, \forall x \in N$ and $\|\xi_0 - \xi\| \leq \delta_\xi K$.

(c) Every densely defined closable derivation on N_0 is inner.

(d) Every closable, conservative, symmetric c.c.n. map on N_0 is bounded in $\|\cdot\|_1$ on $(N_0)_1$.

(e) $\exists F' \subset N_0$ finite and $K' > 0$ such that if $\phi : N \to N$ is a c.p. map with $\phi(1) \leq 1$, $\tau \circ \phi \leq \tau$, $\phi = \phi^*$, and if $\delta'_\phi = \max_{x\in F'}\{\|x - \phi(x)\|_2\}$ then $\tau((y - \phi(y))y^*) \leq K'\delta'_\phi$, $\forall y \in (N)_1$.

Then (b) $\Rightarrow$ (c) $\Rightarrow$ (d) $\Rightarrow$ (e) $\Rightarrow$ (a). If moreover N_0 contains a non-Γ set then we also have (a) $\Rightarrow$ (b).

Proof. (b) $\Rightarrow$ (c). Let $\delta : N_0 \to \mathcal{H}$ be a closable derivation, note that by 1.2 we may assume that δ is real. Let $\phi_t : N \to N$ be the semigroup of normal symmetric c.p. maps associated with δ. Then $\forall y \in N_0$ $\|\delta(y)\|^2 = \lim_{t\to\infty} \tau(\frac{y-\phi_t(y)}{t}y^*)$.

Let $(\mathcal{H}_t, \xi_t)$ be the pointed correspondence obtained from ϕ_t, then since ϕ_t is unital and symmetric $\|y\xi_t - \xi_t y\|_2^2 = 2\tau((y - \phi_t(y))y^*)$ $\forall y \in N$. Let $F \subset N_0$ and $K > 0$ be as in (b) and let $C = \sup_{0<t\leq 1, x\in F} \tau((x - \phi_t(x))x^*)/t$. Then:

$$\|\delta(y)\|^2 = \lim_{t\to 0} \tau(\frac{y - \phi_t(y)}{t}y^*)$$

$$
\begin{aligned}
&= \lim_{t\to 0} \|y\xi_t - \xi_t y\|_2^2/2t \\
&\leq 2 \sup_{0<t\leq 1, x\in F} \|x\xi_t - \xi_t x\|_2^2 K^2/t \\
&= 4 \sup_{0<t\leq 1, x\in F} \tau((x-\phi_t(x))x^*)K^2/t = 4CK^2,
\end{aligned}
$$

for all $y \in (N_0)_1$. Thus by Theorem 2.2 we have that δ is inner.

(c) $\Rightarrow$ (d). This follows from Theorem 2.2.

(d) $\Rightarrow$ (e). Let $\{x_n\}_{n\in\mathbb{N}}$ be a sequence in $(N_0)_1$ such that $N_0 = \text{sp}\{x_n\}_{n\in\mathbb{N}}$. If (e) does not hold then for each $k \in \mathbb{N}$ there exists a c.p. map $\phi_k : N \to N$ such that $\phi_k(1) \leq 1$, $\tau \circ \phi_k \leq \tau$, $\phi_k = \phi_k^*$, and there exists $y_k \in (N_0)_1$ such that $\tau((y_k - \phi_k(y_k))y_k^*) > 4^k \delta'_k$ where $\delta'_k = \max_{j\leq k}\{\|x_j - \phi_k(x_j)\|_2\}$.

Let $\Psi_k = (\text{id} - \phi_k)/\delta'_k$, and let $\Psi = \Sigma_{k=1}^{\infty} 2^{-k}\Psi_k$. Then since $N_0 = \text{sp}\{x_n\}_{n\in\mathbb{N}}$, $\Psi : N_0 \to L^2(N,\tau)$ is a well defined symmetric c.c.n. map with $\Psi(1) \geq 0$. Also since $\phi_k(1) \leq 1$, $\tau \circ \phi_k \leq \tau$, and $\phi_k = \phi_k^*$ if we let $(\mathcal{H}_k, \xi_k)$ be the pointed Hilbert N-N bimodule corresponding to ϕ_k then we have $2\tau((x - \phi_k(x))x^*) \geq \tau \circ \phi_k(x^*x) + \tau(x^*x\phi_k(1)) - 2\tau(\phi(x)x^*) = \|x\xi_k - \xi_k x\|^2 \geq 0$, $\forall x \in N$. Thus

$$
\begin{aligned}
\|\Psi(y_k)\|_1 &\geq \tau(\Psi(y_k)y_k^*) \\
&\geq 2^{-k}\tau((y_k - \phi_k(y_k)y_k^*)/\delta'_k > 2^k,
\end{aligned}
$$

for all $k \in \mathbb{N}$. Hence if we let $\Psi'(x) = \Psi(x) - x\Psi(1)/2 - \Psi(1)x/2$ then Ψ' is a closable, conservative, symmetric c.c.n. map which is unbounded in $\|\cdot\|_1$ on $(N_0)_1$.

(e) $\Rightarrow$ (a). Let F' and K' be as in (e) and let $\varepsilon > 0$. Suppose $\phi : N \to N$ is a c.p. map such that $\phi(1) \leq 1$, $\tau\circ\phi \leq \tau$, $\phi = \phi^*$, and $\|x - \phi(x)\|_2 < \varepsilon^2/2K'$, $\forall x \in F'$. Let $(\mathcal{H}_\phi, \xi_\phi)$ be the pointed Hilbert N-N bimodule associated with ϕ, then since $\|\phi(1)\|_2 \leq 1$ by Lemma 1.1.3 of [P2] we have that $\|y - \phi(y)\|_2^2 \leq \|y\xi_\phi - \xi_\phi y\|^2 = 2\tau((y - \phi(y))y^*) \leq 2K'\delta'_\phi < \varepsilon^2$, $\forall y \in (N)_1$. Hence by Lemma 3 of [PeP] N has property (T).

(a) $\Rightarrow$ (b). If N_0 contains a non-Γ set then Remark 4.1.6 of [P1] shows that one can then apply Proposition 1 of [CJ] to obtain the desired result. □

We note that (d) $\Rightarrow$ (e) in Theorem 3.2 can be suitably adapted to the case of inclusions of σ-compact and locally compact groups thus showing that an inclusion of groups has relative property (T) if and only if "δ depends linearly on ε", answering a question of Jolissaint (see Theorem 1.2 in [J]).

Let $B \subset N$, $1 \in B$ be a $*$-subalgebra and $X = X \in N$. Recall from [V1] that a dual operator to $(X; B)$ in $L^2(N,\tau)$ is an operator $Y \in \mathcal{B}(L^2(N,\tau))$ such that

$$[B, Y] = 0 \ \text{ and } \ [X, Y] = P_1$$

where P_1 is the orthogonal projection onto $\mathbb{C}1$.

3.3. Corollary. *Suppose that N is a separable finite factor with property (T), let $B \subset M$, $1 \in B$ be a $*$-subalgebra and $X = X^* \in N$ such that $B[X]$ generates N as a von Neumann algebra. Suppose that B contains a diffuse element and $B[X]$ contains a non-Γ set, then the conjugate variable $J(X : B)$ does not exist in $L^2(N, \tau)$, i.e. $\Phi^*(X : B) = \infty$. Also $(X; B)$ does not have a dual operator in $L^2(N, \tau)$.*

Proof. If the conjugate variables $J(X : B)$ did exist in $L^2(N, \tau)$ then as in 1.6.1 we would have a closeable derivation on $B[X]$ which is not inner. Therefore by Theorem 3.2 this cannot happen.

The fact that $(X; B)$ does not have a dual system in $L^2(N, \tau)$ then follows directly from [V1]. □

4. Property (T) and amalgamated free products.

We include here an application of the above ideas, showing that a large class of amalgamated free products do not have property (T). We first prove that if N has property (T) then even though a c.c.n. maps may be unbounded on some domains it must still satisfy a certain condition on its rate of growth.

4.1. Theorem. *Suppose N is a finite von Neumann algebra with normal faithful tracial state τ, if N has property (T) and $\Psi : D_\Psi \to L^2(N, \tau) \subset L^1(N, \tau)$ is a conservative, symmetric c.c.n. map, then given any sequence $\{x_n\}_n$ in $(D_\Psi)_1$ such that $\|\Psi(x_n)\|_2 \to \infty$, we have that $\|\Psi(x_n)\|_2 / \|\Psi(x_n)\| \to 0$.*

Proof. Let $\{\Phi_t\}_t$ be the semigroup of unital normal symmetric c.p. maps associated with Ψ as in 1.7, and for each $\beta > 0$ let $\varepsilon_\beta = \sup_{t \leq \beta, x \in N_1} \|\Phi_t(x) - x\|_2$. Since N has property (T) we have that $\varepsilon_\beta \to 0$ as $\beta \to 0$.

For all $\beta > 0$, and $x \in (D_\Psi)_1$ we have:

$$\int_0^\beta \Phi_t \circ \Psi(x) dt = \lim_{s \to 0} \int_0^\beta \Phi_t((\Phi_s(x) - x)/s) dt$$

$$= \lim_{s \to 0} \frac{1}{s} \Big(\int_0^\beta \Phi_{t+s}(x) dt - \int_0^\beta \Phi_t(x) dt \Big)$$

$$= \lim_{s \to 0} \frac{1}{s} \Big(\int_s^{\beta+s} \Phi_t(x) dt - \int_0^\beta \Phi_t(x) dt \Big)$$

$$= \lim_{s \to 0} \frac{1}{s} \Big(\int_\beta^{\beta+s} \Phi_t(x) dt - \int_0^s \Phi_t(x) dt \Big)$$

$$= \Phi_\beta(x) - x.$$

Hence for all $x \in (D_\Psi)_1$:

$$\|\Psi(x)\|_2 \leq \|\frac{1}{\beta}\int_0^\beta \Phi_t \circ \Psi(x)dt\|_2 + \|\frac{1}{\beta}\int_0^\beta (\Phi_t \circ \Psi(x) - \Psi(x))dt\|_2$$

$$\leq \frac{\varepsilon_\beta}{\beta} + \frac{1}{\beta}\int_0^\beta \|(\Phi_t \circ \Psi(x) - \Psi(x))\|_2 dt$$

$$\leq \frac{\varepsilon_\beta}{\beta} + \|\Psi(x)\|\frac{1}{\beta}\int_0^\beta \varepsilon_t dt$$

$$\leq \frac{\varepsilon_\beta}{\beta} + \|\Psi(x)\|\varepsilon_\beta.$$

Thus $\varepsilon_\beta \geq \|\Psi(x)\|_2\beta/(1 + \|\Psi(x)\|\beta)$ and since $\varepsilon_\beta \to 0$ the result follows. □

4.2. Corollary. *Let N_1 and N_2 be finite von Neumann algebras with normal faithful tracial states τ_1 and τ_2 respectively, suppose that B is a common von Neumann subalgebra such that $\tau_1|_B = \tau_2|_B$. Suppose also that there are unitaries $u_i \in \mathcal{U}(N_i)$ such that $E_B(u_i) = 0$, $i = 1, 2$. Then $M = N_1 *_B N_2$ does not have property (T).*

Proof. Let $\tau = \tau_1 *_B \tau_2$ be the trace for M and let $\mathcal{H} = L^2(M,\tau) \otimes_B L^2(M,\tau)$. Define δ to be the unique derivation from the algebraic amalgamated free product to $\mathcal{H}$ which satisfies $\delta(a) = a \otimes_B 1 - 1 \otimes_B a$, $\forall a \in N_1$, and $\delta(b) = 0$, $\forall b \in N_2$. By Corollary 5.4 in [NShSp] $\delta^*(1 \otimes_B 1) = 0$ and so in particular just as in the non-amalgamated case we have that δ is a closable derivation and furthermore if u_1, u_2 are the unitaries as above, and $z \in N_0$ then:

$$\langle \delta((u_1u_2)^n), \delta(z)\rangle = \Sigma_{j=0}^{n-1}\langle (u_1u_2)^j u_1 \otimes_B u_2(u_1u_2)^{n-j-1} - (u_1u_2)^j \otimes_B (u_1u_2)^{n-j}, \delta(z)\rangle$$

$$= \Sigma_{j=0}^{n-1}\langle 1 \otimes_B 1, u_1^*(u_2^*u_1^*)^j\delta(z)(u_2^*u_1^*)^{n-j-1}u_2^* - (u_2^*u_1^*)^j\delta(z)(u_2^*u_1^*)^{n-j}\rangle,$$

also for each $0 \leq j < n$, by using the Leibnitz rule for the derivation we may rewrite $(u_2^*u_1^*)^j\delta(z)(u_2^*u_1^*)^{n-j}$ as a sum of three terms:

$$\delta((u_2^*u_1^*)^j z (u_2^*u_1^*)^{n-j}) \tag{1}$$

$$-\Sigma_{k=0}^{j-1}(u_2^*u_1^*)^k u_2^*\delta(u_1^*)(u_2^*u_1^*)^{j-k-1}z(u_2^*u_1^*)^{n-j} \tag{2}$$

$$-\Sigma_{i=0}^{n-j-1}(u_2^*u_1^*)^j z(u_2^*u_1^*)^i u_2^*\delta(u_1^*)(u_2^*u_1^*)^{n-j-i-1}, \tag{3}$$

however when we take the inner product with $1\otimes_B 1$ the first term will be 0 as mentioned above, and by freeness (since u_1, and u_2 have expectation 0) the other terms will be 0 except when $i = n - j - 1$ in the third term where we have

$$-\langle 1\otimes_B 1, (u_2^*u_1^*)^j z(u_2^*u_1^*)^{n-j-1}u_2^*\delta(u_1^*)\rangle = -\langle 1\otimes_B 1, (u_2^*u_1^*)^j z(u_2^*u_1^*)^{n-j}\otimes_B 1\rangle$$

$$= -\tau(E_B((u_1u_2)^{n-j}z^*(u_1u_2)^j)).$$

Similarly

$$\langle 1\otimes_B 1, u_1^*(u_2^*u_1^*)^j\delta(z)(u_2^*u_1^*)^{n-j-1}u_2^*\rangle = \tau(E_B(u_2(u_1u_2)^{n-j-1}z^*(u_1u_2)^j u_1)).$$

Hence from the above equalities we have:

$$\langle\delta((u_1u_2)^n), \delta(z)\rangle = \Sigma_{j=0}^{n-1}\tau(u_2(u_1u_2)^{n-j-1}z^*(u_1u_2)^j u_1 + (u_1u_2)^{n-j}z^*(u_1u_2)^j)$$

$$= 2n\tau((u_1u_2)^n z^*).$$

In particular if Ψ is the c.c.n. map associated with δ then we have that $\Psi((u_1u_2)^n) = 2n(u_1u_2)^n$ and so $\|\Psi((u_1u_2)^n)\|_2 \to \infty$ but $\|\Psi((u_1u_2)^n)\|_2/\|\Psi((u_1u_2)^n)\| \not\to 0$, hence by Theorem 4.1 M does not have property (T). □

Note that the only place where we used the fact that u_1 and u_2 were unitairies was to insure that $2n\|(u_1u_2)^n\|_2 \to \infty$. Also note that the conditions of the above Corollary are satisfied when M is a free product (with amalgamation over $\mathbb{C}$) as well as when M is a group von Neumann algebra coming from an amalgamated free products of groups. We also mention that from the calculation above we are able to compute explicitly the semigroup of c.p. maps that δ generates, it is the semigroup given by $\phi_t = (e^{-2t}\mathrm{id} + (1 - e^{-2t})E_B) *_B \mathrm{id}$.

Acknowledgments. The author would like to thank Professor Sorin Popa for his encouragement and many stimulating discussions. The author would also like to thank Professor Dima Shlyakhtenko for showing him the references [S1] and [S2], and Professor Jean-Luc Sauvageot for his useful comments on an earlier version of this paper.

References

[AW] C.A. Akemann, M.E. Walter: *Unbounded negative definite functions*, Can, J. Math. **33** (1981), 862-871.

[BdHV] B. Bekka, P. de la Harpe, A. Valette: *Kazhdan's property (T)*, Book Preprint, (July 2002).

[ChE] E. Christensen, D.E. Evans: *Cohomology of operator algebras and quantum dynamical semigroups*, J. London Math. Soc. **20** (1979), 358-368.

[CiSa] F. Cipriani, J.-L. Sauvageot: *Derivations as square roots of Dirichlet forms*, J. Funct. Anal. **201** (2003), no. 1, 78-120.

[C1] A. Connes: *Classification of injective factors*, Ann. of Math. **104** (1976), 73-115.

[C2] A. Connes: *Classification des facteurs*, Proc. Symp. Pure Math. **38** (Amer. Math. Soc., 1982), 43-109.

[C3] A. Connes: *Correspondences*, hand-written notes, 1980.

[CJ] A. Connes, V.F.R. Jones: *Property (T) for von Neumann algebras*, Bull. London Math. Soc. **17** (1985), 57-62.

[DL] E. B. Davies, J. M. Lindsay: *Non-commutative symmetric Markov semigroups*, Math. Z. **210** (1992), 379-411.

[De] P. Delorme: 1-*cohomologie des représentations unitaires de groupes de Lie semisimples et résolubles. Produits tensoriels continus et représentations*, Bull. Soc. Math. France, **105** (1977), 281-336.

[Ef] E. G. Effros: *Property* Γ *and inner amenability*, Proc. Amer. Math. Soc. **47** (1975), 483-486.

[E] D.E. Evans: *Conditionally completely positive maps on operator algebras*, Quart. J. Math. Oxford **4** (1977), 271-284.

[G] A. Guichardet: *Etude de la* 1-*cohomologie et de la topologie du dual pour les groupes de Lie à radical abélien*, Math. Ann., **228** (1977), 215-232.

[J] P. Jolissaint: *On Property (T) for Pairs of Topological Groups*, preprint 2004.

[Ka] D.A. Kazhdan: *Connection of the dual space of a group with the structure of its closed subgroups*, Funct. Anal. Appl. **1** (1967), 63-65.

[Ki] A. Kishimoto: *Dissipations and derivations*, Commun. math. Phys. **47** (1976) 25-32.

[L] G. Lindblad: *On the generators of quantum dynamical semigroups*, Commun. math. Phys. **48** (1976), 119-130.

[MvN] F. J. Murray, J. von Neumann: *On rings of operators* IV, Ann. of Math. (2) **44** (1943), 716-808.

[NShSp] A. Nica, D. Shlyakhtenko, R. Speicher: *Operator valued distributions. I. Characterizations of freeness*, Int. Math. Res. Not. (2002), 1509-1538.

[PeP] J. Peterson, S. Popa: *On the notion of relative property (T) for inclusions of von Neumann algebras*, J. Funct. Anal., **219** (2005), no. 2, 469-483.

[P1] S. Popa: *Correspondences*, INCREST preprint 1986, unpublished.

[P2] S. Popa: *On a class of type* II_1 *factors with Betti numbers invariants*, preprint math.OA/0209130, to appear in Ann. of Math.

[S1] J.-L. Sauvageot: *Tangent bimodules and locality for dissipative operators on C*-algebras*, Quantum probability and appl., IV, Lecture notes in Math. **1396** (1989), 322-338.

[S2] J.-L. Sauvageot: *Quantum Dirichlet forms, differential calculus and semigroups*, Quantum probability and appl., V, Lecture notes in Math. **1442** (1990), 334-346.

[V1] D. Voiculescu: *The analogues of entropy and of Fisher's information measure in free probability theory* V. *Noncommutative Hilbert transform*, Invent. math. **132** (1998) 189-227.

[V2] D. Voiculescu: *The analogues of entropy and of Fisher's information measure in free probability theory* VI. *Liberation and mutual free information*, Adv. Math. **146** (1999) 101-166.

MATH DEPT UCLA, LOS ANGELES, CA 90095-155505
E-mail address: `jpete@math.ucla.edu`